

UNDERUSE AND MISUSE OF DATA FROM BEACHED BIRD SURVEYS

FRANCIS K. WIESE¹ & KIM ELMSLIE²

¹*North Pacific Research Board, 1007 West 3rd Avenue, Suite 100, Anchorage, Alaska, 99501, USA
(francis.wiese@nprb.org)*

²*International Fund for Animal Welfare, 1 Nicholas Street, Suite 612, Ottawa, Ontario, K1N 7B7, Canada*

Received 22 March 2006, accepted 27 October 2006

WIESE, F.K. & ELMSLIE K. 2006. Underuse and misuse of data from beached bird surveys. *Marine Ornithology* 34: 157–159.

INTRODUCTION

Beached bird surveys continue to be one of the main mechanisms by which marine oil pollution is monitored, and one of the earliest signals that an oiling event, small or large, has taken place. Beached bird surveys have been or are currently conducted in a number of countries around the world (Canada east coast: Wiese & Ryan 2003, Campbell 2005; Canada west coast: Burger 1993, O'Hara & Morgan 2006; USA west coast: Speich & Wahl 1986, Bodkin & Jameson 1991, Nur *et al.* 1997, Lyday *et al.* In press; USA east coast: Simons 1985; Argentina: Perkins 1983, Boersma 1995, Harris *et al.* 2006; Japan: Ohata *et al.* 1993; New Zealand: Veitch 1982; Australia: Raaymakers 1995; South Africa: Avery 1989; North Sea: Dahlmann *et al.* 1994; UK: Cadbury 1978, Stowe & Underwood 1984, Heubeck 1995; Shetland Islands: Heubeck 2006; Denmark: Joensen & Hansen 1977; Belgium: Kuyken 1978; Netherlands: Camphuysen 1989, 1998; Germany: Averbek *et al.* 1992; France: Debout 1984, Raavel 1990; Lithuania: Vaitakus *et al.* 1994, Žydelis & Dagys 1997, Žydelis *et al.* 2006). Despite regional successes in reducing chronic oil pollution (e.g. Camphuysen 1998, Camphuysen & Heubeck 2001), oiled birds continue to wash ashore, and many unmonitored regions around the globe likely face similar problems.

Beached bird surveys are conducted either to help document wildlife mortality after a large oiling event or as part of regular (monthly, weekly) monitoring programs to document trends in chronic oil pollution. The first objective is reactive and restricted in time and space; the timing is crucial for achieving valid mortality estimates and overall damage assessments. The second objective can be both proactive and reactive. It is long-term, and it requires a rigorous and systematic approach for the duration of the program. Although the immediate goal of surveys is to collect data on beach conditions and bird casualties, one overall aim should be to quantify the problem in ways that help to increase awareness and raise regional, national, and international standards for dealing with marine oil pollution. That broader objective is where programs often fall short. Most of them quantify the number of birds found per kilometer of coastline and the proportion of birds found that are oiled ("oiling rate"), they generally do not advance knowledge any further than that. Thus beached bird surveys remain primarily a monitoring tool, when they could also be used as a tool for change.

A TOOL FOR CHANGE

To make beached bird surveys a tool for change rather than just a monitoring exercise, the data collected should be augmented in three important ways:

- Determine factors responsible for trends in the data, or at least establish whether a given trend reflects a change in clean or oiled

birds. See Camphuysen (1989, 1998), Wiese & Ryan (2003), and Wiese & Robertson (2004) for a discussion of factors such as wind, temperature, distribution, and hunting pressure that influence the numbers of clean and oiled birds found on beaches.

- Determine as accurately as possible the total number of birds affected—managers and policymakers want an extrapolated number, not just an oiling rate or the number of birds found. Without an overall mortality estimate, the true scope of the problem will never be confronted (see Wiese & Robertson 2004 on determining seabird mortality from chronic pollution). If accurate estimates can be derived, then valid assessments of spatial and temporal trends in mortality and contributing factors can also be determined, permitting comparisons between various regions of the world.
- Determine the impact of oil pollution on seabird populations—a final step that is difficult to achieve and roundly debated even after large, well-documented spills. Unequivocally linking oil pollution and fluctuations in population sizes is difficult because of uncertainties in mortality estimates, a dearth of baseline data, imprecision in censusing seabird colonies, the existence of other anthropogenic pressures occurring simultaneously and natural variation (Wiese *et al.* 2004). Still, the need for accurate mortality estimates and assessment of population-level impacts cannot be overstated.

Unfortunately, these steps—particularly the latter two—are missing in most places, a situation that has long been a major hindrance to policy change. As noted, the problem is measured regionally, but it is global in scale. If adequate efforts were made to determine oiled seabird mortality in places where rigorous beached bird surveys are now conducted, the ability of scientists to influence the global policies of the International Maritime Organization (IMO), and consequently of the countries signatory to the International Convention for the Prevention of Pollution from Ships (MARPOL), would be strengthened significantly.

UNDERUSE AND MISUSE OF SURVEY DATA

To achieve the identified goals, more data need to be collected and fuller advantage needs to be taken of data collected independently that can inform the patterns determined from current scientific observations.

For example, information on birds should include morphometrics, age, sex, freshness (scavenging rate), cause of death, persistence and detection probabilities. Factors influencing deposition, such as wind, tides and currents, can generally be quantified using data from outside sources (e.g. weather service, universities). Combined with information about the distribution and abundance of birds at sea, such data help to elucidate factors influencing mortality levels and trends. Past studies incorporating this approach effectively focused

on temperature, wind, oceanography, shipping (number of vessels transiting an area), fishing activities and hunting (e.g. Kampp 1991, Camphuysen 1998, Chardine 1998, Wiese *et al.* 2004).

Clearly, many of the tools necessary to achieve reduced chronic oil pollution and fewer seabirds affected exist, and yet they are often not fully applied. Still worse are instances in which data are misused or misinterpreted, either because of a limited knowledge of the physical and biologic processes involved, or willfully because of political motivations. Thus, reports on the total numbers of birds encountered, numbers of birds found per kilometer or the number of oil slicks detected, including trends in data that were collected with inconsistent methodology, have all been used to advance political agendas. Deliberations generally occur in the absence of oversight by either the public or the scientists who collected the data. It is thus the responsibility of scientists to present their information in a manner that precludes misinterpretation, by incorporating all available evidence to clarify and corroborate observed patterns and trends.

OBSTACLES TO CHANGE

Paradoxically, science itself can be an obstacle to changing public policy, and misuse of scientific results by individuals unable or unmotivated to interpret the data correctly should be anticipated. Other obstacles include limited public awareness, political interests and legislative restrictions. Those issues are largely outside the realm of science, but scientists can no longer afford to ignore them. It is vital that scientists understand where their information is being communicated and how to increase the effectiveness of the message. By becoming an integral part of awareness campaigns—even by lobbying—scientists can enhance the credibility of their message and help to guard against misinterpretation. Such involvement need not sacrifice scientific integrity or objectivity, and its pursuit is necessary in today's political climate.

LEGAL REGIMES

Many laws and agreements govern how nations deal with the deliberate dumping of oil and the occurrence of oiled seabirds. In Canada, these instruments include MARPOL, the *Canada Shipping Act*, the *Canadian Environmental Protection Act*, the *Migratory Bird Convention Act*, the *Fisheries Act*, the *Species at Risk Act*, the *Oceans Act* and the Offshore Petroleum Boards.

Most prosecutions of oil polluters in Canada are carried out under the *Canada Shipping Act*, which deals primarily with poor shipping practices. It does not support direct charges because of affected wildlife or deleterious effects on the marine environment, as might occur with some of the other acts mentioned. There is, however, no reason that an offender cannot be charged under several acts simultaneously, as has been successful in many cases in the United States (e.g. the *Anax Liberian* in 1998). Effective methods of deterrence for marine polluters include, besides high prosecution rates, the presence of aerial surveillance, detention of ships and substantial fines.

In Atlantic Canada, little aerial surveillance has been conducted in relation to a vast area of jurisdiction, vessels have only rarely been directed to port and detained and, until recently, imposed fines have been low by international standards, making them effective deterrents to pollution. Each of these shortcomings has contributed to a failure to stop or significantly curtail the deliberate dumping of oil in Canadian waters.

Fortunately, some of this changed for Canada when Bill C-15 (www.ec.gc.ca/press/2005/050519_n_e.htm), containing amendments to the Migratory Birds Convention Act and the Canadian Environmental Protection Act, was passed on 19 May 2005. Successful passage of the bill is a credit to the many individuals and organizations throughout Canada who worked for decades on the issue of chronic oil pollution. Now, more than 50 years after the problem of oiled seabirds on Newfoundland beaches was identified (Tuck 1961), the law supports enforcement out to the 200-mile limit; increases funds for aerial surveillance; makes it possible to arrest and charge captains, crew and owners of polluting vessels; and imposes higher fines.

How the new law will be implemented and how effective it will be in reducing seabird deaths attributable to the deliberate dumping of oil off the Newfoundland coast from the currently estimated level of 300 000 annually (Wiese & Robertson 2004) are open questions that only time will answer. In any event, good science was one essential factor in this victory for animal welfare and environmental health. Among other studies, beached bird surveys were the foundation of that science, and thus that activity has indeed become a tool for change.

CONCLUSIONS

Because oiled beached birds are often the only indication that oil has been spilled at sea, it is crucial that those beachings be well documented and correctly interpreted. Beach surveys can help to ensure that national and international policymakers deal successfully with the global problem of oil dumping. Agencies responsible for enforcing anti-dumping laws have a natural tendency to use any downward fluctuations in the reported numbers of oiled birds as evidence of job performance. Others argue that the numbers are meaningless if not placed in an environmental context that specifies when, where and how birds are deposited on beaches. The discussion ranges from biology and methods—species vulnerabilities, weather regimes, currents and survey designs—to political and economic considerations such as MARPOL and the cost of transportation. All play a role in how policymakers choose to interpret the results of beached bird surveys. It is the job of the scientific community to acquire and present the relevant data in such a manner that the principal outcome is heightened public awareness and the political will to appropriately deal with chronic marine oil pollution.

ACKNOWLEDGMENTS

We thank all the dedicated individuals at Environment Canada, the Canadian Coast Guard, Memorial University of Newfoundland, the International Fund for Animal Welfare, Nature Canada and the World Wildlife Fund who, through their dedication, continue to improve the health of Canada's marine ecosystems.

REFERENCES

- AVERBECK, C., KORSCH, M. & VAUK, G. 1992. Der Einfluss von Oelverschmutzungen auf Seevoegel an den deutschen Nordseekuesten von 1984–1990. *Seevoegel* 13: 12–16.
- AVERY, G. 1989. Results of patrols for beached seabirds conducted in southern Africa in 1984 and 1985. *Cormorant* 17: 57–71.
- BODKIN, J.L. & JAMESON, R.J. 1991. Patterns of seabird and marine mammal carcass deposition along the central Californian coast, 1980–1986. *Canadian Journal of Zoology* 69: 1149–1155.

- BOERSMA, P.D. 1995. Chronic oil pollution is a large source of mortality for Magellanic Penguins (*Spheniscus magellanicus*). *Pacific Seabirds* 22: 26.
- BURGER, A.E. 1993. Mortality of seabirds assessed from beached-bird surveys in southern British Columbia. *Canadian Field Naturalist* 107: 164–176.
- CADBURY, C.J. 1978. The beached bird survey and other seabird surveillance. *Ibis* 120: 119–120.
- CAMPBELL, G. 2005. Cape Breton beached bird survey: four-year report, 2001–2005. Bird Studies Canada Atlantic Region Internal Report, November 21, 2005. Sackville, NB: Environmental Canada. 15 pp.
- CAMPHUYSEN, C.J., 1998. Beached bird surveys indicate decline in chronic oil pollution in the North Sea. *Marine Pollution Bulletin* 36: 519–526.
- CAMPHUYSEN, C.J. 1989. Beached bird surveys in the Netherlands 1915–1988: seabird mortality in the southern North Sea since the early days of oil pollution. Technisch Rapport Vogelbescherming 1. Amsterdam: Werkgroep Noordzee. 308 pp.
- CAMPHUYSEN, C.J. & HEUBECK, M. 2001. Marine oil pollution and beached bird surveys: the development of a sensitive monitoring instrument. *Environmental Pollution* 112: 443–461.
- CHARDINE, J.W. 1998. Review of the seabird bycatch problem in Arctic Canada. In: Baken, V. & Falk, K. (Eds). Incidental take of seabirds in commercial fisheries in the Arctic countries. *Circumpolar Seabird Working Group Technical Report* 1: 9–14.
- DAHLMANN, G., TIMM, D., AVERBECK, C., CAMPHUYSEN, C., SKOV, H. & DURINCK, J. 1994. Oiled seabirds—comparative investigations on oiled seabirds and oiled beaches in the Netherlands, Denmark and Germany (1990–93). *Marine Pollution Bulletin* 28: 305–310.
- DEBOUT, G. 1984. Les décomptes d’oiseaux échoués: 1974–1982. *Le Cormoran* (Special issue): 1–3.
- HARRIS, R.J., TSENG, F.S., POKRAS, M.A., SUEDEMEYER, B.A., BOGART, J.S.H., PRESCOTT, R.L. & NEWMAN, S.H. 2006. Seabird Ecological Assessment Network (SEANET) volunteer beached bird surveys in Massachusetts, 2003–2004. *Marine Ornithology* 34: 115–122.
- HEUBECK, M. 1995. Shetland beached bird surveys: national and European context. *Proceedings of the Royal Society of Edinburgh* 103B: 165–179.
- HEUBECK, M. 2006. The Shetland beached bird survey, 1979–2004. *Marine Ornithology* 34: 123–127.
- JOENSEN, A.H. & HANSEN, E.B. 1977. Oil pollution and seabirds in Denmark 1971–1976. *Danish Review of Game Biology* 10: 1–31.
- KAMPP, K. 1991. Mortality of Thick-billed Murres in Greenland inferred from band recovery data. *Canadian Wildlife Service Occasional Papers* 69: 15–22.
- KUYKEN, E. 1978. Beached bird surveys in Belgium. *Ibis* 120: 122–123.
- LYDAY, S., ROLETTTO, J., MORTENSON, J. & HALL, J. In press. Beach Watch: sanctuary ecosystem assessment surveys. *Pacific Seabirds*.
- NUR, N., SYDEMANN, W.J., PYLE, P., STENZEL, L.E., AINLEY, D.G. & SCHUSTER, T.G. 1997. Temporal, spatial and species specific patterns of chronic oil pollution as revealed by the beached bird surveys, Farallon oiled bird survey, and bird rescue programs in Central California [Unpublished report]. Sacramento, CA: California Department of Fish and Game, Office of Oil Spill Prevention and Response. 64 pp.
- O’HARA, P. & MORGAN, K.H. 2006. Do low rates of oiled seabird carcass recoveries in beached bird surveys indicate low rates of ship-source oil spills? *Marine Ornithology* 34: 133–140.
- OHATA, K., NEMOTO, S. & MURAI, M. 1993. Mortality of seabirds due to oil contamination along the Tomakomai coast, Hokkaido. *Strix* 12: 214–218.
- PERKINS, J. 1983. Oiled Magellanic Penguins in Golfo San Jose, Argentina. *Marine Pollution Bulletin* 14: 383–387.
- RAAYMAKERS, S. 1995. Illegal “operational” discharges continue in Barrier Reef. *Marine Pollution Bulletin* 30: 362.
- RAEVEL, P. 1990. Bilan de 20 années de recensement des oiseaux morts sur le littoral du Nord Pas-de-Calais à la fin de l’hiver: exemple de l’intérêt d’une enquête à protocole déterminé. *Le Heron* 23: 159–167.
- SIMONS, M.M. Jr. 1985. Beached bird survey project on the Atlantic Gulf Coasts. *American Birds* 39: 358–362.
- SPEICH, S.M. & WAHL, T.R. 1986. Rates of occurrence of dead birds in Washington’s inland marine waters, 1978 and 1979. *Murrelet* 67: 51–59.
- STOWE, T.J. & UNDERWOOD, L.A. 1984. Oil spillages affecting seabirds in the United Kingdom, 1966–1983. *Marine Pollution Bulletin* 15: 147–152.
- TUCK, L.M. 1961. The murre. Canadian Wildlife Monograph Series 1. Ottawa: Queen’s Printer. 260 pp.
- VAITAKUS, G., PETRAITIS, A. & ŽYDELIS, R. 1994. Beached bird density trends in Lithuania during 1991–1994. *Acta Ornithologica Lithuanica* 9–10: 78–86.
- VEITCH, C.R. 1982. Seabirds found dead in New Zealand in 1980. *Notornis* 29: 41–47.
- WIESE, F.K. & ROBERTSON, G.J. 2004. Assessing impacts of chronic oil discharges at sea on seabirds: a general oiled seabird mortality model applied to Eastern Canada. *Journal of Wildlife Management* 68: 627–638.
- WIESE, F.K., ROBERTSON, G.J. & GASTON, A.J. 2004. Impacts of chronic marine oil pollution and the murre hunt in Newfoundland on Thick-billed Murre populations in the Eastern Canadian Arctic. *Biological Conservation* 116: 205–216.
- WIESE, F.K. & RYAN, P.C. 2003. The extent of chronic marine oil pollution in southeastern Newfoundland waters assessed through beached-bird surveys 1984–1999. *Marine Pollution Bulletin* 46: 1090–1101.
- ŽYDELIS, R. & DAGYS, M. 1997. Winter period ornithological impact assessments of oil related activities and sea transportation in Lithuanian inshore waters of the Baltic Sea and in the Kurši Lagoon. *Acta Zoologica Lithuanica* 6: 45–65.
- ŽYDELIS, R., DAGYS, M. & VAITKUS, G. 2006. Beached bird surveys in Lithuania, Eastern Baltic: reflection of marine oil pollution and bird mortality in fishing nets. *Marine Ornithology* 34: 161–166.