NUMBER AND DISTRIBUTION RECORDS OF CASPIAN TERN HYDROPROGNE CASPIA IN INDONESIA, WITH SPECIAL REFERENCE TO SUMATRA

MUHAMMAD IQBAL¹, YUS RUSILA NOOR², DENI MULYANA³, HENNI MARTINI⁴, FERRY HASUDUNGAN⁵, IWAN FEBRIANTO⁶, FRANSISCA NONI TIRTANINGTYAS⁷, ARUM SETIAWAN⁸, INDRA YUSTIAN⁸ & HILDA ZULKIFLI⁸

¹Biology Programe, Faculty of Science, Sriwijaya University, Jalan Padang Selasa 524, Palembang 30139, Indonesia (kpbsos26@yahoo.com)

²Yayasan Lahan Basah (Wetlands International Indonesia), JalanBango No.11, Tanah Sareal, Bogor, West Java, Indonesia

³Berbak Sembilang National Park, South Sumatra office, Jalan Tanjung Api-api komplek Imadinatuna No. 114, South Sumatra, Indonesia

⁴Hutan Kita Institute (HAKI), Jalan Yudo No. 9 H, Palembang, South Sumatra 30126, Indonesia

⁵Ecology Conservation of Nature and Wildlife in Indonesia Foundation (EKSAI), Jalan Kutisari 1 No. 19, Surabaya, East Java, Indonesia

⁶Burung Indonesia, Jalan Dadali No. 32, Tanah Sereal, Bogor, East Java, Indonesia

⁷Burung Laut Indonesia, Depok, East Java 16421, Indonesia ⁸Department of Biology, Faculty of Science, Sriwijaya University, Jalan Raya Palembang-Prabumulih km 32, Indralaya, Sumatera Selatan 30662, Indonesia

Received 01 June 2020, accepted 04 August 2020

ABSTRACT

IQBAL, M., NOOR, Y.R., MULYANA, D., MARTINI, H., HASUDUNGAN, F., FEBRIANTO, I., TIRTANINGTYAS, F.N., SETIAWAN, A., YUSTIAN, I. & ZULKIFLI, H. 2020. Number and distribution records of Caspian Tern *Hydroprogne caspia* in Indonesia, with special reference to Sumatra. *Marine Ornithology* 48: 297–301.

We summarize all reports of Caspian Tern *Hydroprogne caspia* recorded in Indonesia, with particular focus on Sumatra (1983–2019). There were 29 records in total, which together indicate that the Caspian Tern is a regular visitor to Indonesia, in contrast to previous assumptions that it is rare. This species has been recorded during the non-breeding period (September to March in Indonesia, and August to April in Sumatra) in small numbers ranging from one to 40 birds. Many potential sites for this species, particularly along east coast of Sumatra, have still not been well explored. Additional search efforts will advance our knowledge of Caspian Tern and other seabirds that occur in Indonesia.

Key words: Caspian Tern, historical records, Indonesia, Sumatra

Caspian Tern *Hydroprogne caspia* is a monotypic, cosmopolitan species occurring in North America, North Europe, Africa, Madagascar, Asia, Australia, and New Zealand (del Hoyo & Collar 2014, Gochfeld & Burger 1996). In the Northern Hemisphere, it winters in the Caribbean, and on the coasts of Africa, the Indian Ocean, southern China, and Southeast Asia to the Thai-Malay Peninsula and Sumatra (Harrison 1983, Wells 1999). In Australia, the Caspian Tern breeding season occurs March–November (Higgins & Davies 1996, Dunlop & McNeill); in New Zealand, birds breed October–December (Barlow & Dowding 2002). The global population is estimated at 250 000–470 000 individuals (Birdlife International 2020), and the Southeast Asian population is estimated at 10 000–25 000 birds (Delany & Scott2006).

In Indonesia, the Caspian Tern is a rare, mostly coastal migrant, but it is occasionally present within inland wetlands (Eaton *et al.* 2016). The species is a vagrant in Greater Sundas, recorded rarely on the coast of east Sumatra and in Sabah, Sarawak, and Brunei (MacKinnon & Phillipps 1993, Mann 2008). Van Marle & Voous (1988) stated that Caspian Tern is a winter visitor in Sumatra. It has been reported in Wallacea, at Timor, in October and January, possibly having migrated from Australia (White & Bruce 1986). Caspian Tern is a scarce visitor in New Guinea, mainly to the coastal Trans-Fly Region, but it is not clear whether this species occurs in Indonesian Papua (west New Guinea) (Pratt & Beehler 2015, Gregory 2017).

Caspian Tern is a highly distinctive bird, and any records that appear to be ambiguous or unconfirmed should be rejected. Based on the ease of its identification, we compiled and summarized all published and unpublished records of the species in Sumatra and Indonesia, including information found on social media and from internet sources, e.g., Facebook groups of local birdwatchers, eBird (eBird 2017), Global Biodiversity Information Facility (GBIF 2020), and iNaturalist (iNaturalist 2020). We screened all information for authenticity and correct species identification.

Since they were first reported in Jambi province in 1983, 29 records of Caspian Tern have accumulated in Indonesia and Sumatra (Table 1). There are eight records of Caspian Tern observed outside Sumatra Island. In Sumatra, the locations where this species has been observed can be divided into three major areas along the east coast: Percut (North Sumatra, *ca.* 03°43′N, 098°46′E), Cemara beach and Berbak (Jambi province, *ca.* 01°28′S, 104°20′E), and the Banyuasin Peninsula (South Sumatra province, *ca.* 02°14′S, 104°50′E). No inland records have been reported (Fig. 1). The number of Caspian Terns per record in Sumatra is relatively small, in the range of 1–40 birds. One Caspian Tern was ringed with the Sumatra flag on 17 January 2010 at Cemara Beach (Noni & Londo 2010), but it has not been resighted.

Many species of terns are migratory, breeding in the extreme northern and southern latitudes and only frequenting the tropics in




Fig. 1. Locations of Caspian Tern recorded in Indonesia and Sumatra. Numbers refer to adjacent yellow circles: (1) Percut and adjacent areas, North Sumatra province; (2) Cemara beach and Berbak, Jambi province; (3) Banyuasin peninsula, South Sumatra province; (4) Maros, South Sulawesi province; (5) Losari beach, South Sulawesi province; (6) Makasar old harbour, South Sulawesi province; (7) Sanur beach, Bali province; (8) Bipolo, East Nusa Tenggara province; (9) Kupang, East Nusa Tenggara province; (10) Sorong, West Papua province; and (11) Aru island, Maluku province.

their respective winter seasons (MacKinnon & Phillipps 1993). In the Malay Peninsula, the Caspian Tern is a locally common non-breeding visitor occurring in small numbers on the west coast and south to Singapore, where it is rare (Wells 1999, Jeyarajasingam & Pearson 2012). Its presence has been reported at 15 sites in the Malay Peninsula (eBird 2017), with the largest number—100 birds—observed at the Kapar Power Station, Selangor (Craig 2017). In Singapore, two observations of one single bird were reported at two locations (eBird 2017).

A summary of records indicate that Caspian Tern is a regular non-breeding visitor, occurring in small numbers, in Indonesia and Sumatra (Table 1, Fig. 2). Records from Indonesia and Sumatra span September–March and August–April, respectively. In Sumatra, most records occur in October (5), followed by December (4); other months have one to two records (August, September, November, January, and April have two records; February has one record). No records of Caspian Tern were reported in Indonesia and Sumatra from May–July or during the Northern Hemisphere summer breeding season.

Accurate spatial information on Caspian Tern and other seabirds in Indonesia are needed to assess population trends in this region. An increasing number of birdwatchers in Indonesia and Sumatra, supported by high magnification optics, has led to increasing documentation of vagrant and migratory seabirds during the last decade, which has complicated accurate assessment of population dynamics (Iqbal & Albayquni 2016, Hasyim *et al.* 2019, Tirtaningtyas & Yordan 2017, Iqbal *et al.* 2020, Setiawan *et al.* 2020). Many potential sites along the east coast of Sumatra have not been rigorously explored, particularly in Aceh, Riau, Bangka-Belitung, and Lampung province. More extensive, quantified search efforts of coastal seabirds around the mangrove beaches of these areas is required to advance our knowledge of Caspian Tern and other seabirds in Sumatra.

ACKNOWLEDGEMENTS

We thank Andrew Crossland, Chairunas Adha Putra, and James Eaton for sharing their information on Caspian Tern in Sumatra. We are very grateful to the editors of *Marine Ornithology*, and anonymous reviewers for critical review of the draft.

No.	Date	No. of Caspian Terns recorded	Sites	Sources
1	14 Oct 1983	2	Cemara beach, Jambi province	van Marle & Voous 1988, Silvius et al. 1986
2	27 Oct 1984	1	Cemara beach, Jambi province	van Marle & Voous 1988, Silvius et al. 1986
3	28 Oct 1984	3	Cemara beach (near Labuhan Pering), Jambi province	van Marle & Voous 1988, Silvius et al. 1986
4	24 Nov 1984	8	Cemara beach (near Tanjung Jabung), Jambi province	van Marle & Voous 1988, Silvius et al. 1986
5	Feb-Mar 1985	Present, no count	Kupang bay, East Nusa Tenggara province	Trainor & Hidayat 2014
6	Aug-Oct 1985	Present, no count	Kupang bay, East Nusa Tenggara province	Trainor & Hidayat 2014
7	02 Mar 1988	1 (possibly this species)	Aru island, Maluku, Maluku province	Bishop 2017
8	01 Dec 1988	11	Banyuasin Peninsula, South Sumatra province	Verheugt et al. 1993, Holmes 1996
9	01 Aug 1989	2	Banyuasin Peninsula, South Sumatra province	Verheugt et al. 1993, Holmes 1996
10	23-30 Dec 1995	6	Percut, North Sumatra province	Crossland et al. 2012, van Balen et al. 2013
11	15–17 April 2001	Present, no count	Jambat river, Berbak, near Cemara beach, Jambi province	Wetland International Indonesia Programme 2001
12	30 Oct 2002	3	Cemara beach, Jambi province	Hasudungan & Wardoyo 2002
13	31 Aug 2003	2	Dinding river, Banyuasin Peninsula, South Sumatra province	Iqbal 2002
14	28 Apr 2006	1	Banyuasin Peninsula, South Sumatra province	MI pers. obs
15	22 Dec2009	1	Cemara beach, Jambi province	FNT & IF pers. obs
16	17 Jan 2010	1	Cemara beach, Jambi province	Noni & Londo 2010
17	02 Oct 2010	1	Datuk Alam beach, near Percut, North Sumatra province	Andrew Crossland pers. comm.
18	04 Sept 2012	1	Sejara beach, near Percut, North Sumatra province	Andrew Crossland pers. comm.
19	28 Jan 2017	40	Banyuasin Peninsula, South Sumatra province	YRN pers. obs.
20	08 Sept 2017	6	Banyuasin Peninsula, South Sumatra province	MI, DM & HM pers. obs.
21	12 Sept 2017	2	Maros, South Sulawesi province	Upton 2017
22	02 Oct 2017	2	Losari beach, South Sulawesi	Evans et al. 2017
23	20 Oct 2017	Present, no count	Makasar old harbour, South Sulawesi province	Buskirik 2017
24	20 Feb2018	6	Banyuasin Peninsula, South Sumatra province	MI, DM & HM pers. obs
25	04 Dec 2018	4	Sanur beach, Bali province	Gardner & Gardner 2017
26	04 Feb 2019	2	Sorong, West Papua province	Grabin & Grabin 2017
27	04 Sept 2019	1	Bipolo, East Nusa Tenggara province	Ovin et al. 2019
28	20-30 Nov 2019	15	Cemara beach, Jambi province	IF pers. obs.
29	21 Dec 2019	20	Banyuasin Peninsula, South Sumatra province	MI & DM pers. obs.

Marine Ornithology 48: 297-301 (2020)




Fig. 2. Caspian Terns found in Sumatra, Indonesia. (A) Two terns encountered among shorebirds on 21 December 2019 along the Banyuasin Peninsula, Banyuasin district, South Sumatra province; (B) tern observed in flight on 22 December 2009 at Cemara Beach, Tanjung Jabung Timur district, Jambi province; (C) terns observed on 28 January 2017 along the Banyuasin Peninsula, Banyuasin district, South Sumatra province. Photos taken by Muhammad Iqbal [A], Fransisco Noni Tirtaningtyas [B], and Yus Rusila Noor [C].

REFERENCES

- BARLOW, M.L. & DOWDING, J.E. 2002. Breeding biology of Caspian terns (*Sterna caspia*) at a colony near Invercargill, New Zealand. *Notornis* 49: 76–90.
- BIRDLIFE INTERNATIONAL 2020. Species factsheet: Hydroprogne caspia. Cambridge, UK: BirdLife International. [Accessed online at http://www.birdlife.org on 01 June 2020.]
- BISHOP, D. 2017. *Checklist S17989957*. Ithaca, USA: eBird. [Accessed online at https://ebird.org/checklist/S17989957 on 27 June 2020.]
- BUSKIRIK, W. 2017. *Checklist S39929291*. Ithaca, USA: eBird. [Accessed online at https://ebird.org/checklist/S39929291on 27 June 2020.]
- CHANDLER, R. 2009. Shorebirds of the Northern Hemisphere. London, UK: Christopher Helm.
- CRAIG, M. 2017. *Kapar Power Station*. Ithaca, USA: eBird. [Accessed online at https://ebird.org/hotspot/L1122479 on 27 June 2020.]
- CROSSLAND, A.C., LUBIS, L., SINAMBELA, S.A., SITORUS, A.S., SITORUS, A.W. & MUIS, A. 2012. Observations of shorebirds along the Deli-Serdang coast, North Sumatra province, Indonesia: 1995–2006. *Stilt* 61: 37–44.
- DEL HOYO, J. & COLLAR, N. J. 2014. HBW and BirdLife International Illustrated Checklist of the Birds of the World, 1. Barcelona, Spain: Lynx Edicions.
- DELANY, S. & SCOTT, D. 2006. *Waterbird Population Estimates*. Wageningen, Netherlands: Wetlands International.
- DUNLOP, J.N. & MCNEILL, S. 2017. Local movements, foraging patterns, and heavy metals exposure in Caspian Terns *Hydroprogne caspia* breeding on Penguin Island, Western Australia. *Marine Ornithology* 45: 115–120.

- EATON, J.A., VAN BALEN, B., BRICKLE, N.W. & RHEINDT, F.E. 2016. *Birds of the Indonesian Archipelago*. Barcelona, Spain: Lynx Edicions.
- EBIRD. 2017. *eBird: An online database of bird distribution and abundance* [web application]. Ithaca, USA: eBird. [Accessed online at http://www.ebird.org on 27 June 2020.]
- EVANS, B., DOHERTY, D. & EVANS, T. 2017. *Checklist S45172317*. Ithaca, USA: eBird. [Accessed online at https://ebird.org/checklist/S45172317 on 27 June 2020.]
- GARDNER, V. & GARDNER, M. 2017. *Checklist S50464471*. Ithaca, USA: eBird. [Accessed online at https://ebird.org/australia/checklist/S50464471 on 27 June 2020].
- GBIF (GLOBAL BIODIVERSITY INFORMATION FACILITY) 2020. *Hydroprogne caspia* (Pallas, 1770). [Accessed online at https://www.gbif.org/species/2481236 on 27 June 2020.]
- GOCHFELD, M. & BURGER, J. 1996. Family Sternidae (Terns).
 In: J. DEL HOYO, A. ELLIOT & J. SARGATAL (Eds).
 Handbook of the Birds of the World. Vol. 3. Hoatzin to Auks, pp. 624-667. Barcelona, Spain: Lynx Edicions.
- GRABIN, M. & GRABIN, B. 2017. Checklist S68342324. Ithaca, USA: eBird. [Accessed online at https://ebird.org/me/checklist/ S68342324 on 27 June 2020.]
- GREGORY, P. 2017. Birds of New Guinea, Including Bismarck Archipelago and Bougainville. Barcelona, Spain: Lynx Edicions.
- HARRISON, P. 1983. Seabirds, an Identification Guide. Boston, USA: Houghton Mifflin Company.
- HASUDUNGAN, F. & WARDOYO, A. 2002. *Pemantauan Kawasan Berbak No. 4, Okt-Nov 2002. Laporan Teknis Dokumen Proyek No. 64.* Palembang, Indonesia: Proyek Konservasi Terpadu Lahan Basah Pesisir Berbak Sembilang. [In Indonesian]
- HASYIM, A., IQBAL, M., SETIAWAN, A. & YUSTIAN, A. 2019. Status of Black-Headed Gull *Larus ridibundus* in Indonesian Borneo. *Marine Ornithology* 47: 215–216.
- HIGGINS, P.J. & DAVIES, S.J.J.F. 1996. Handbook of Australian, New Zealand 6 Antarctic Birds. Vol. 3. Snipe to Pigeons. Melbourne, Australia: Oxford University Press.
- HOLMES, D.A.1996. Sumatra bird report. Kukila 8: 9-56.
- INATURALIST 2020. Caspian Tern (Hydroprogne caspia).
 Berkeley, USA: iNaturalist. [Accessed online at https://www.inaturalist.org/guide_taxa/393825 on 01 June 2020.]
- IQBAL, M. & ALBAYQUNI, A.A. 2016. First record of a Slaty-backed gull *Larus schistisagus* for indonesia. *Marine Ornithology* 44: 135–136.
- IQBAL, M. 2002. Pemantauan Kawasan Sembilang No. 7, Juli/ Agustus 2003. Laporan Teknis No. 74. Palembang, Indonesia: Proyek Konservasi Terpadu Lahan Basah Pesisir Berbak Sembilang. [In Indonesian]
- IQBAL, M., ABDILLAH, H., FEBRIANTO, I., AMRUL, H.M., WINDUSARI, Y. & HANUM, L. 2020. Recent status of Blackheaded Gull *Chroicocephalus ridibundus* in Sumatra, Indonesia. *Marine Ornithology* 48: 53–54.
- JAYARAJASINGAM, A. & PEARSON, A. 2012. A field guide to the birds of Peninsular Malaysia and Singapore. Oxford, UK: Oxford University Press.
- MACKINNON, J. & PHILLIPPS, K. 1993. A Field Guide to the Birds of Borneo, Sumatra, Java and Bali. Oxford, UK: Oxford University Press.
- MANN, C.F. 2008. *The Birds of Borneo*. Peterborough, UK: British Ornithologists' Union (Check-list 23).
- MCCLURE, H.E. 1998. *Migration and Survival of the Birds of Asia*. Bangkok, Thailand: White Lotus Co., Ltd.

- NONI, F. & LONDO, I.2010. Field report for Wildlife Conservation Society, Sumatra, Indonesia: December 2009–January 2010. *Tattler* 18: 8–10.
- OLSEN, K.M. & LARSSON, H. 2003. *Gulls of Europe, Asia and North America*. London, UK: Christopher Helm.
- OVIN, P., BATES, D. & SMITH, J. 2019. *Checklist S59966910*. Ithaca, USA: eBird. [Accessed online at https://ebird.org/checklist/S59966910 on 27 June 2020.]
- PRATT, T.K. & BEEHLER, B.M. 2015. *Birds of New Guinea*. Princeton, USA: Princeton University Press.
- SETIAWAN, A., IQBAL, M., PORMANSYAH, YUSTIAN, I. & ZULKIFLI, H. 2020. Recent records of Little Cormorant *Microcarbo niger* in Sumatra, Indonesia. *Marine Ornithology* 48: 161–162.
- SILVIUS, M.J. & VERHEUGHT, W.J.M. 1986. The birds of Berbak Game Reserve, Jambi province, Sumatra. *Kukila* 2: 76–84.
- SONOBE, K. & USUI, S. 1993. A Field Guide to the Waterbirds of Asia. Tokyo, Japan: Wild Bird Society of Japan.
- TIRTANINGTYAS, F.N. & YORDAN, K. 2017. Updating the seabird fauna of Jakarta Bay, Indonesia. *Marine Ornithology* 45: 11–16

- TRAINOR C.R. & HIDAYAT, O. 2014. Kupang Bay: an internationally significant wetland in West Timor, Indonesia. BirdingAsia 21: 45–50
- UPTON, N. 2017. Checklist S39379192. Ithaca, USA: eBird. [Accessed online at https://ebird.org/checklist/S39379192 on 27 June 2020.]
- VAN BALEN, B., TRAINOR, C. & NOSKE, R. 2013. Around the archipelago. *Kukila* 17: 41–72.
- VAN MARLE, J.G. & VOOUS, K.H. 1988. *The Birds of Sumatra:* an Annotated Check-list. Tring, UK: British Ornithologists' Union (Check-list 10).
- VERHEUGT, W.J.M., SKOV, H. & DANIELSEN, F. 1993. Notes on the birds of the tidal lowlands and floodplains of South Sumatra province, Indonesia. *Kukila* 6: 53–84.
- WELLS, D.R. 1999. *Birds of the Thai–Malayan Peninsula*. Vol. 1. London, UK: Academic Press.
- WETLAND INTERNATIONAL INDONESIA PROGRAMME. 2001. Survei Pengkajian Cepat Taman Nasional Berbak, Jambi. Palembang, Indonesia: Proyek Konservasi Terpadu Lahan Basah Pesisir Berbak Sembilang. [In Indonesian]
- WHITE, C.M.N. & BRUCE, M.D. 1986. *The Birds of Wallacea* (Sulawesi, The Moluccas & Lesser Sunda Islands, Indonesia). Tring, UK: British Ornithologists' Union (Check-list 7).